

Vaccine Administration

Vaccine Administration

Pages 1-12

Resources

- Needle Length
- Comforting Restraint
- Giving All the Doses
- After the Shots
- Medical Management of Vaccine Reaction



Vaccine Administration

Appropriate vaccine administration is critical to vaccine effectiveness. The recommended site, route and dosage for each vaccine are based on clinical trials, practical experience and theoretical considerations. The following information provides general guidelines for administration of vaccines for those who administer vaccines, as well as those in training, education and supervisory positions. This information should be used in conjunction with professional standards for medication administration, vaccine manufacturers' product guidelines, CDC's Advisory Committee on Immunization Practices (ACIP) General Recommendations on Immunization, the American Academy of Pediatrics' (AAP) Report of the Committee on Infectious Diseases Red Book, and state/agency-related policies and procedures. An education plan that includes competency-based training on vaccine administration should be considered for all persons who administer vaccines to children or adults.

Preparation

Patient Preparation - Patients should be prepared for vaccination with consideration for their age and stage of development. Parents/guardians and patients should be encouraged to take an active role before, during and after the administration of vaccines.

- **Screening** - All patients should be screened for contraindications and precautions for each scheduled vaccine. Many state immunization programs and other organizations have developed and made available standardized screening tools. Sample screening forms for children and adults are available from the Immunization Action Coalition (www.immunize.org).
- **Vaccine Safety & Risk Communication** - Parents/guardians and patients are exposed through the media to information about vaccines, some of which is inaccurate or misleading. Healthcare providers should be prepared to discuss the benefits and risks of vaccines using Vaccine Information Statements (VIS) and other reliable resources. Establishing an open dialogue provides a safe, trust-building environment in which individuals can freely evaluate information, discuss vaccine concerns and make informed decisions regarding immunization.
- **Atraumatic Care** - Vaccine safety issues and the need for multiple injections have increased the concerns and anxiety associated with immunizations. Healthcare providers need to display confidence and establish an environment that promotes a sense of security and trust for the patient and family, utilizing a variety of techniques to minimize the stress and discomfort associated with receiving injections. This is particularly important when administering vaccines to children.

- **Positioning & Comforting Restraint** - The healthcare provider must accommodate for the patient's comfort, safety, age, activity level and the site of administration when considering patient positioning and restraint. For a child, the parent/guardian should be encouraged to hold the child during administration. If the parent is uncomfortable, another person may assist or the patient may be positioned safely on an examination table.
- **Pain Control** - Pain is a subjective phenomenon influenced by multiple factors, including an individual's age, anxiety level, previous healthcare experiences and culture. Consideration for these factors is important as the provider develops a planned approach to management of injection pain.
 - **Topical Anesthetics or a Vapocoolant Spray** - May be applied to decrease pain at the injection site. These products should be used only for the ages recommended and as directed by the product manufacturer.
 - **Analgesic Agents** - A non-aspirin-containing pain reliever may be considered to decrease discomfort and fever following vaccination. These products should be used only in age-appropriate doses.
 - **Diversiónary Techniques** - Age-appropriate non-pharmacologic techniques may provide distraction from pain associated with injections. Diversion can be accomplished through a variety of techniques.
 - **Dual Administrators** - Some providers favor the technique of two individuals simultaneously administering vaccines at separate sites. The premise is that this procedure may decrease anxiety from anticipation of the next injection(s). The effectiveness of this procedure in decreasing pain or stress associated with vaccine injections has not been evaluated.

Infection Control - Healthcare providers should follow Standard Precautions to minimize the risks of spreading disease during vaccine administration.

- **Hand-washing** - The single, most effective disease prevention activity is good hand-washing. Hands should be washed thoroughly with soap and water, or cleansed with an alcohol-based waterless antiseptic between patients, before vaccine preparation or any time hands become soiled (e.g., diapering, cleaning, etc.).
- **Gloves** – Unless dictated by clinic policies, gloves are not required to be worn when administering vaccines unless the person administering the vaccine is likely to come into contact with potentially infectious body fluids or has open lesions on the hands. It is important to remember that gloves cannot prevent needle stick injuries.
- **Needle Stick Injuries** - Should be reported immediately to the site supervisor, with appropriate care and follow-up given as directed by state/local guidelines. Safety needles or needle-free injection devices should be used if available to reduce the risk of injury.



- **Equipment Disposal** - Used needles should not be detached from syringes, recapped or cut before disposal. All used syringe/needle devices should be placed in puncture proof containers to prevent accidental needle sticks and reuse. Empty or expired vaccine vials are considered medical waste and should be disposed of according to state and local regulations.

Vaccine Preparation - Proper vaccine handling and preparation is critical in maintaining the integrity of the vaccine during transfer from the manufacturer's vial to the syringe and ultimately to the patient.

- **Equipment Selection**

- **Syringe Selection** - A separate needle and syringe should be used for each injection. A parenteral vaccine may be delivered in either a 1-mL or 3-mL syringe as long as the prescribed dosage is delivered. Syringe devices with sharps engineered sharps injury protection are available, recommended by OSHA, and required in many states to reduce the incidence of needle stick injuries and potential disease transmission. Personnel should be involved in evaluation and selection of these products. Staff should receive training with these devices before using them in the clinical area.

- **Needle Selection** - Vaccine must reach the desired tissue site for optimal immune response. Therefore, needle selection should be based upon the prescribed route, size of the individual, volume and viscosity of the vaccine, and injection technique. (See Subcutaneous & Intramuscular Injections, below.) Typically, vaccines are not highly viscous, and therefore a fine gauge needle (22-25 gauge) can be used.

- **Needle-Free Injection** - A new generation of needle-free vaccine delivery devices has been developed in an effort to decrease the risks of needle stick injuries to healthcare workers and to prevent improper reuse of syringes and needles. For more information on needle-free injection technology, see the CDC website:
www.cdc.gov/vaccines/dev/jetinject.htm.

- **Inspecting Vaccine** - Each vaccine vial should be carefully inspected for damage or contamination prior to use. The expiration date printed on the vial or box should be checked. Vaccine can be used through the last day of the month indicated by the expiration date unless otherwise stated on the package labeling. Expired vaccine should never be used.

- **Reconstitution** - Some vaccines are prepared in a lyophilized form that requires reconstitution, which should be done according to manufacturer guidelines. Diluent solutions vary; use only the specific diluent supplied for the vaccine. Once reconstituted, the vaccine must be either administered within the time guidelines provided by the manufacturer or discarded.



Changing the needle after reconstitution of the vaccine is not necessary unless the needle has become contaminated or bent. Continue with standard medication preparation guidelines.

- **Prefilling Syringes** - CDC strongly discourages filling syringes in advance, because of the increased risk of administration errors. Once the vaccine is in the syringe, it is difficult to identify the type or brand of vaccine. Other problems associated with this practice are vaccine wastage and possible bacterial growth in vaccines that do not contain a preservative. Furthermore, medication administration guidelines state that the individual who administers a medication should be the one to draw up and prepare it. An alternative to prefiling syringes is to use filled syringes supplied by the vaccine manufacturer. Syringes other than those filled by the manufacturer are designed for immediate administration, not for vaccine storage.

In certain circumstances, such as a large influenza clinic, more than one syringe can be filled. One person should prefill only a few syringes at a time, and the same person should administer them. Any syringes left at the end of the clinic day should be discarded.

Under no circumstances should MMR, varicella or zoster vaccines ever be reconstituted and drawn prior to the immediate need for them. These live virus vaccines are unstable and begin to deteriorate as soon as they are reconstituted with diluent.


- **Labeling** - Once a vaccine is drawn into a syringe, the content should be indicated on the syringe. There are a variety of methods for identifying or labeling syringes (e.g., keep syringes with the appropriate vaccine vials, place the syringes in a labeled partitioned tray, or use color-coded labels or preprinted labels).

Administration

Route - Administering a vaccine by the recommended route is imperative. Deviation from the recommended route of administration might reduce vaccine efficacy or increase the risk of local reactions.


Subcutaneous (Sub-Q or SC)

Subcutaneous injections are administered into the fatty tissue found below the dermis and above muscle tissue.



Lynne Larson, www.biovisuals.com

Subcutaneous Site - Subcutaneous tissue can be found all over the body. The usual sites for vaccine administration are the thigh (for infants <12 months of age) and the upper outer triceps of the arm (for persons >12 months of age). If necessary, the upper outer triceps area can be used to administer subcutaneous injections to infants.




Subcutaneous Needle Gauge & Length - 5/8-inch, 23- to 25-gauge needle

Subcutaneous Technique - Follow standard medication administration guidelines for site assessment/selection and site preparation. To avoid reaching the muscle, pinch up the fatty tissue, insert the needle at a 45° angle and inject the vaccine into the tissue. Withdraw the needle and apply light pressure to the injection site for several seconds with a dry cotton ball or gauze.




Intramuscular (IM)

Intramuscular injections are administered into muscle tissue below the dermis and subcutaneous tissue.




Lynne Larson, www.biovisuals.com


Intramuscular Site - Although there are several IM injection sites on the body, the recommended IM sites for vaccine administration are the vastus lateralis muscle (anterolateral thigh) and the deltoid muscle (upper arm). The site depends on the age of the individual and the degree of muscle development.



The vastus lateralis muscle of the upper thigh used for intramuscular injections.



The vastus lateralis site of the right thigh, used for an intramuscular injection.





Intramuscular Needle Gauge - 22- to 25-gauge needle

Intramuscular Needle Length - For all intramuscular injections, the needle should be long enough to reach the muscle mass and prevent vaccine from seeping into subcutaneous tissue, but not so long as to involve underlying nerves, blood vessels, or bone. The vaccinator should be familiar with the anatomy of the area into which the vaccine will be injected. Decision on needle size and site of injection must be made for each person on the basis of the size of the muscle, the thickness of adipose tissue at the injection site, the volume of the material to be administered, injection technique, and the depth below the muscle surface into which the material is to be injected.

Infants (Younger Than 12 Months)

For the majority of infants, the anterolateral aspect of the thigh is the recommended site for injection because it provides a large muscle mass. The muscles of the buttock have not been used for administration of vaccines in infants and children because of concern about potential injury to the sciatic nerve, which is well documented after injection of antimicrobial agents into the buttock. If the gluteal muscle must be used, care should be taken to define the anatomic landmarks. If the gluteal muscle is chosen, injection should be administered lateral and superior to a line between the posterior superior iliac spine and the greater trochanter or in the ventrogluteal site, the center of a triangle bounded by the anterior superior iliac spine, the tubercle of the iliac crest, and the upper border of the greater trochanter.

Injection technique is the most important factor to ensure efficient intramuscular vaccine delivery. If the subcutaneous and muscle tissue are bunched to minimize the chance of striking bone, a 1-inch needle is required to ensure intramuscular administration in infants. For the majority of infants, a 1-inch, 22-25-gauge needle is sufficient to penetrate muscle in an infant's thigh. For newborn (first 28 days of life) and premature infants, a 5/8-inch needle usually is adequate if the skin is stretched flat between thumb and forefinger and the needle inserted at a 90-degree angle to the skin.

Toddlers and Older Children (12 Months through 10 Years)

The deltoid muscle should be used if the muscle mass is adequate. The needle size for deltoid site injections can range from 22 to 25 gauge and from 5/8 to 1 inch on the basis of the size of the muscle and the thickness of adipose tissue at the injection site. A 5/8-inch needle is adequate only for the deltoid muscle and only if the skin is stretched flat between thumb and forefinger and the needle inserted at a 90° angle to the skin. For toddlers, the anterolateral thigh can be used, but the needle should be at least 1 inch in length.



Adolescents and Adults (11 Years or Older)

For adults and adolescents, the deltoid muscle is recommended for routine intramuscular vaccinations. The anterolateral thigh also can be used. For men and women weighing less than 130 lbs (60 kg) a 5/8-1-inch needle is sufficient to ensure intramuscular injection. For women weighing 130-200 lbs (60-90 kg) and men 130-260 lbs (60-118kg), a 1-1½-inch needle is needed. For women weighing more than 200 lbs (90 kg) or men weighing more than 260 lbs (118 kg), a 1½-inch needle is required.

Technique - Follow standard medication administration guidelines for site assessment/selection and site preparation. To avoid injection into subcutaneous tissue, spread the skin of the selected vaccine administration site taut between the thumb and forefinger, isolating the muscle. Another technique, acceptable mostly for pediatric and geriatric patients, is to grasp the tissue and “bunch up” the muscle. Insert the needle fully into the muscle at a 90° angle and inject the vaccine into the tissue. Withdraw the needle and apply light pressure to the injection site for several seconds with a dry cotton ball or gauze.

- **Aspiration** - Aspiration is the process of pulling back on the plunger of the syringe prior to injection to ensure that the medication is not injected into a blood vessel. Although this practice is advocated by some experts, the procedure is not required because no large blood vessels exist at the recommended injection sites.
- **Multiple Vaccinations** - When administering multiple vaccines, NEVER mix vaccines in the same syringe unless approved for mixing by the Food and Drug Administration (FDA). If more than one vaccine must be administered in the same limb, the injection sites should be separated by 1-2 inches so that any local reactions can be differentiated. Vaccine doses range from 0.2 mL to 1 mL. The recommended maximum volume of medication for an IM site, varies among references and depends on the muscle mass of the individual. However, administering two IM vaccines into the same muscle would not exceed any suggested volume ranges for either the vastus lateralis or the deltoid muscle in any age group. The option to also administer a subcutaneous vaccine into the same limb, if necessary, is acceptable since a different tissue site is involved.
- If a vaccine and an immune globulin preparation are administered simultaneously (e.g., Td/Tdap and tetanus immune globulin [TIG] or hepatitis B vaccine and hepatitis B immune globulin [HBIG]), a separate anatomic site should be used for each injection. The location of each injection should be documented in the patient’s medical record.

- **Nonstandard Administration** - Deviation from the recommended route, site and dosage of vaccine is strongly discouraged and can result in inadequate protection. In situations where nonstandard administration has occurred, refer to the ACIP General Recommendation on Immunization, *MMWR* 2006; 55 (RR-15), for specific guidance.

Special Situations

Bleeding Disorders - Individuals with a bleeding disorder or who are receiving anticoagulant therapy may develop hematomas in IM injection sites. Prior to administration of IM vaccines the patient or family should be instructed about the risk of hematoma formation from the injection. Additionally, a physician familiar with the patient's bleeding disorder or therapy should be consulted regarding the safety of administration by this route. If the patient periodically receives hemophilia replacement factor or other similar therapy, IM vaccine administration ideally should be scheduled shortly after replacement therapy. A 23-gauge or finer needle should be used and firm pressure applied to the site for at least two minutes. The site should not be rubbed or massaged.

Latex Allergy - Administration of a vaccine supplied in a vial or syringe that contains natural rubber (refer to product information) should not be administered to an individual with a history of a severe (anaphylactic) allergy to latex, unless the benefit of vaccination clearly outweighs the risk of an allergic reaction. These situations are rare. Medical consultation and direction should be sought regarding vaccination. A local or contact sensitivity to latex is not a contraindication to vaccination.

Syncopal or Vasovagal Response - This "fainting" may occur during vaccine administration, especially with adolescents and adults. Because individuals may fall and sustain injury as a result, the provider should have the patient sit during injection(s). A syncopal or vasovagal response is not common and is not an allergic reaction. However, if syncope develops, the provider should observe and administer supportive care until the patient is recovered.

Anaphylaxis (a life-threatening acute allergic reaction) - Each facility that administers vaccines should have a protocol, procedures and equipment to provide initial care for suspected anaphylaxis. Facility staff should be prepared to recognize and respond appropriately to this type of emergency situation. All staff should maintain current CPR certification. Emergency protocols, procedures and equipment/supplies should be reviewed periodically. For additional information on medical management of vaccine reactions in children, teens, and adults, see the 2006 ACIP General Recommendations on Immunization (p. 19), the 2006 AAP *Red Book* (pp. 64-66). Although both fainting and allergic reactions are rare, vaccine providers should strongly consider observing patients for 15 minutes after they are vaccinated.

Documentation

All vaccines administered should be fully documented in the patient's permanent medical record. Documentation should include:

1. Date of administration
2. Name or common abbreviation of vaccine
3. Vaccine lot number
4. Vaccine manufacturer
5. Administration site
6. Vaccine Information Statement (VIS) edition date (found in the lower right corner of the back of the VIS)
7. Name and address of vaccine administrator. This should be the address where the record is kept. If immunizations are given in a shopping mall, for example, the address would be the clinic where the permanent record will reside.
8. Nevada Vaccine Program eligibility must be documented if administering a Nevada Vaccine Program supplied vaccine.


Facilities that administer vaccines are encouraged to participate in state/local immunization information systems. The patient or parent should be provided with an immunization record that includes the vaccines administered with dates of administration.

Administering Vaccines: Dose, Route, Site, and Needle Size

Vaccines	Dose	Route
Diphtheria, Tetanus, Pertussis (DTaP, DT, Tdap, Td)	0.5 mL	IM
<i>Haemophilus influenzae</i> type b (Hib)	0.5 mL	IM
Hepatitis A (HepA)	≤18 yrs: 0.5 mL ≥19 yrs: 1.0 mL	IM
Hepatitis B (HepB)	≤19 yrs: 0.5 mL* ≥20 yrs: 1.0 mL	IM
<i>*Persons 11–15 yrs may be given Recombivax HB® (Merck) 1.0 mL adult formulation on a 2-dose schedule.</i>		
Human papillomavirus (HPV)	0.5 mL	IM
Influenza, live attenuated (LAIV)	0.5 mL	Intranasal spray
Influenza, trivalent inactivated (TIV)	6-35 mos: 0.25 mL ≥3 yrs: 0.5 mL	IM
Measles, mumps, rubella (MMR)	0.5 mL	SC
Meningococcal, conjugated (MCV4)	0.5 mL	IM
Meningococcal, polysaccharide (MPSV4)	0.5 mL	SC
Pneumococcal conjugate (PCV)	0.5 mL	IM
Pneumococcal polysaccharide (PPV)	0.5 mL	IM or SC
Polio, inactivated (IPV)	0.5 mL	IM or SC
Rotavirus (Rv)	2.0 mL	Oral
Varicella (Var)	0.5 mL	SC
Zoster (Zos)	0.65 mL	SC
Combination Vaccines		
DTaP+HepB+IPV (Pediarix™) DTaP+Hib (Trihibit™) Hib+HepB (Comvax™)	0.5 mL	IM
MMR+Var (ProQuad®)	≤12 yrs: 0.5 mL	SC
HepA+HepB (Twinrix®)	≥18 yrs: 1.0 mL	IM

Injection Site and Needle Size		
Subcutaneous (SC) injection		
Use a 23–25 gauge needle. Choose the injection site that is appropriate to the person's age and body mass.		
Age	Needle Length	Injection Site
Infants (1–12 mos)	5/8"	Fatty tissue over anterolateral thigh muscle
Children (≥12 mos), adolescents, & adults	5/8"	Fatty tissue over anterolateral thigh muscle or fatty tissue over triceps
Intramuscular (IM) injection		
Use a 22–25 gauge needle. Choose the injection site and needle length appropriate to the person's age and body mass.		
Age	Needle Length	Injection Site
Newborn (1 st 28 days)	5/8"*	Anterolateral thigh muscle
Infant (1–12 mos)	1"	Anterolateral thigh muscle
Toddler (1–2 yrs)	1"–1 1/4" 5/8"*–1"	Anterolateral thigh muscle or deltoid muscle of arm
Children 3–18 yrs	5/8"*–1" 1"–1 1/4"	Deltoid muscle of arm or anterolateral thigh muscle
≥19 yrs (Sex/Weight)		
Male/Female less than 130 lbs	5/8"*–1"	Deltoid muscle of arm
Female (130–200 lbs) Male (130–260 lbs)	1"–1 1/2"	Deltoid muscle of arm
Female (200+ lbs) Male (260+ lbs)	1 1/2"	Deltoid muscle of arm

**If skin is stretched tight and subcutaneous tissue is not bunched.*



Please note: Always refer to the package insert included with each biologic for complete vaccine administration information. CDC's Advisory Committee on Immunization Practices (ACIP) recommendations for the particular vaccine should be reviewed as well.


COMFORTING RESTRAINT

FOR IMMUNIZATIONS

• The method:

This method involves the parent in embracing the child and controlling all four limbs. It avoids “holding down” or overpowering the child, but it helps you steady and control the limb of the injection site.


• For infants and toddlers:



Have parent hold the child on parent's lap.

1. One of the child's arms embraces the parent's back and is held under the parent's arm.
2. The other arm is controlled by the parent's arm and hand. For infants, the parent can control both arms with one hand.
3. Both legs are anchored with the child's feet held firmly between the parent's thighs, and controlled by the parent's other arm.

• For kindergarten and older children:




Hold the child on parent's lap or have the child stand in front of the seated parent.

1. Parent's arms embrace the child during the process.
2. Both legs are firmly between parent's legs.



Giving All the Doses

One way to give 5 doses at one visit



PCV7 (IM)

Hep B (IM)

IPV (SC)

Hib (IM)

DTaP (IM)

(SC injections can also be given in the thigh)

- Injection sites should be separated by 1 inch
- IM injections not recommended in infant's arms

Giving All the Doses

One way to give 7 doses (if needed) at one visit



IPV (SC)

MMR (SC)

DTaP (IM)


Hib (IM)

Varicella (SC)

PCV7 (IM)

HepB (IM)

The deltoid muscle is an option for IM injections in children 18 months and older with adequate muscle mass



Vaccinations may hurt a little . . .
but disease can hurt a lot!

Call your clinic right away if you answer "yes" to any of the following questions:

- Does your child have a temperature about which your health care provider has told you to be concerned?
- Is your child pale or limp?
- Has your child been crying for more than 3 hours and just won't quit?
- Does your child have a strange cry that isn't normal (a high-pitched cry)?
- Is your child's body shaking, twitching, or jerking?
- Does your child have marked decrease in activity or decrease in responsiveness?

Check the back of this page for information on the proper dosage of medication you can give your child to reduce pain or fever.

After the Shots . . .

What to do if your child has discomfort

Your child may need extra love and care after getting vaccinated. Some vaccinations that protect children from serious diseases also can cause discomfort for a while. Here are answers to questions many parents have after their children have been vaccinated. If this sheet doesn't answer your questions, call your clinic or health care provider.

Clinic or health care provider phone number: _____

I think my child has a fever. What should I do?

Check your child's temperature to find out if there is a fever. Do not use a mercury thermometer. If your child is younger than 3 years of age, taking a temperature with a rectal digital thermometer provides the best reading. Once your child is 4 or 5 years of age, you may prefer taking a temperature by mouth with an oral digital thermometer. Tympanic thermometers, which measure temperature inside the ear, are another option for older babies and children. If your child is older than 3 months of age, you can also take an underarm (axillary) temperature, although it is not as accurate.

Here are some things you can do to help reduce fever:

- Give your child plenty to drink.
- Clothe your child lightly. Do not cover or wrap your child tightly.
- Give your child a fever-reducing medication such as acetaminophen (e.g., Tylenol®) or ibuprofen (e.g., Advil®, Motrin®). **Do not give aspirin.** Recheck your child's temperature after 1 hour.
- Sponge your child in 1–2 inches of lukewarm water.
- If your child's temperature is _____°F (_____°C) or higher or, if you have questions, call your clinic or health care provider.

My child has been fussy since getting vaccinated. What should I do?

After vaccination, children may be fussy due to pain or fever. You may want to give your child a medication such as acetaminophen (e.g., Tylenol®) or ibuprofen (e.g., Advil®, Motrin®) to reduce pain and fever. **Do not give aspirin.** If your child is fussy for more than 24 hours, call your clinic or health care provider.

My child's leg or arm is swollen, hot, and red. What should I do?

- Apply a clean, cool, wet washcloth over the sore area for comfort.
- For pain, give a medication such as acetaminophen (e.g., Tylenol®) or ibuprofen (e.g., Advil®, Motrin®). **Do not give aspirin.**
- If the redness or tenderness increases after 24 hours, call your clinic or health care provider.

My child seems really sick. Should I call my health care provider?

If you are worried **at all** about how your child looks or feels, call your clinic or health care provider!

Medications and Dosages to Reduce Pain and Fever

Important notes:

1. Ask your health care provider or pharmacist which formulation is best for your child.
2. Give dose based on your child's weight. If you don't know the weight, give dose based on your child's age. Do not give more medication than recommended.
3. If you have questions about dosing or any other concern, call your clinic or health care provider.
4. Always use a proper measuring device. For example:



- When giving infant drops, use only the dosing device (dropper or syringe) enclosed in the package.





- When giving children's suspension or liquid, use the dosage cup enclosed in the package. If you misplace the dosage cup, consult your health care provider or pharmacist for advice. (Kitchen spoons are not accurate measures.)

5. **WARNING:** If you're also giving your child over-the-counter (OTC) medications such as cold preparations, be aware that these may contain pain or fever reducers such as acetaminophen or ibuprofen. Be sure to read all OTC medication labels carefully to ensure your child is not receiving more acetaminophen or ibuprofen than recommended.

Acetaminophen Dosing Information (Tylenol® or another brand)



Give every 4–6 hours, as needed, no more than 5 times in 24 hours (unless directed to do otherwise by your health care provider).

Weight of child	Age of child	Infant drops  0.8 mL = 80 mg	Children's liquid or suspension  1 tsp (5 mL) = 160 mg	Children's tablets 1 tablet = 80 mg	Junior strength 1 tablet = 160 mg
6–11 lbs (2.7–5 kg)	0–3 mos	Advised dose*: _____			
12–17 lbs (5.5–7.7 kg)	4–11 mos	Advised dose*: _____	Advised dose*: _____		
18–23 lbs (8.2–10.5 kg)	12–23 mos	Advised dose*: _____	Advised dose*: _____		
24–35 lbs (10.9–15.9 kg)	2–3 yrs	1.6 mL	1 teaspoon (160 mg)	2 tablets	
36–47 lbs (16.4–21.4 kg)	4–5 yrs		1½ teaspoons (240 mg)	3 tablets	
48–59 lbs (21.8–26.8 kg)	6–8 yrs		2 teaspoons (320 mg)	4 tablets	2 tablets
60–71 lbs (27.3–32.3 kg)	9–10 yrs		2½ teaspoons (400 mg)	5 tablets	2½ tablets
72–95 lbs (32.7–43.2 kg)	11 yrs		3 teaspoons (480 mg)	6 tablets	3 tablets

*Ask your health care provider

Ibuprofen Dosing Information (Advil®, Motrin® or another brand)

Give every 6–8 hours, as needed, no more than 4 times in 24 hours (unless directed to do otherwise by your health care provider).

Weight of child	Age of child	Infant drops  1.25 mL = 50 mg	Children's liquid or suspension  1 tsp (5 mL) = 100 mg	Children's tablets 1 tablet = 50 mg	Junior strength 1 tablet = 100 mg
under 11 lbs (5 kg)	under 6 mos	Advised dose*: _____			
12–17 lbs (5.5–7.7 kg)	6–11 mos	1.25 mL			
18–23 lbs (8.2–10.5 kg)	12–23 mos	1.875 mL			
24–35 lbs (10.9–15.9 kg)	2–3 yrs		1 teaspoon (100 mg)	2 tablets	
36–47 lbs (16.4–21.4 kg)	4–5 yrs		1½ teaspoons (150 mg)	3 tablets	
48–59 lbs (21.8–26.8 kg)	6–8 yrs		2 teaspoons (200 mg)	4 tablets	2 tablets
60–71 lbs (27.3–32.3 kg)	9–10 yrs		2½ teaspoons (250 mg)	5 tablets	2½ tablets
72–95 lbs (32.7–43.2 kg)	11 yrs		3 teaspoons (300 mg)	6 tablets	3 tablets

*Ask your health care provider

Medical Management of Vaccine Reactions in Children and Teens

All vaccines have the potential to cause an adverse reaction. To minimize adverse reactions, patients should be carefully screened for precautions and contraindications before vaccine is administered. Even with careful screening, reactions can occur. These reactions can vary from trivial and inconvenient (e.g., soreness, itching) to severe and life threatening (e.g., anaphylaxis). If reactions occur, staff should be prepared with procedures for their management. The table below describes procedures to follow if various reactions occur.

Reaction	Symptoms	Management
Localized	Soreness, redness, itching, or swelling at the injection site	Apply a cold compress to the injection site. Consider giving an analgesic (pain reliever) or antipruritic (anti-itch) medication.
	Slight bleeding	Apply an adhesive compress over the injection site.
	Continuous bleeding	Place thick layer of gauze pads over site and maintain direct and firm pressure; raise the bleeding injection site (e.g., arm) above the level of the patient's heart.
Psychological fright and syncope (fainting)	Fright before injection is given	Have patient sit or lie down for the vaccination.
	Extreme paleness, sweating, coldness of the hands and feet, nausea, light-headedness, dizziness, weakness, or visual disturbances	Have patient lie flat or sit with head between knees for several minutes. Loosen any tight clothing and maintain an open airway. Apply cool, damp cloths to patient's face and neck.
	Fall, without loss of consciousness	Examine the patient to determine if injury is present before attempting to move the patient. Place patient flat on back with feet elevated.
	Loss of consciousness	Check the patient to determine if injury is present before attempting to move the patient. Place patient flat on back with feet elevated. Call 911 if patient does not recover immediately.
Anaphylaxis	Sudden or gradual onset of generalized itching, erythema (redness), or urticaria (hives); angioedema (swelling of the lips, face, or throat); severe bronchospasm (wheezing); shortness of breath; shock; abdominal cramping; or cardiovascular collapse	See "Emergency Medical Protocol for Management of Anaphylactic Reactions in Children and Teens" on the next page for detailed steps to follow in treating anaphylaxis.

Supplies Needed

- Aqueous epinephrine 1:1000 dilution, in ampules, vials of solution, or prefilled syringes, including epinephrine auto-injectors (e.g., EpiPen). If EpiPens are to be stocked, both EpiPen Jr. (0.15 mg) and adult EpiPens (0.30 mg) should be available.
- Diphenhydramine (Benadryl) injectable (50 mg/mL solution) and oral (12.5 mg/5 mL suspension) and 25 mg or 50 mg capsules or tablets
- Syringes: 1–3 cc, 22–25g, 1", 1½", and 2" needles for epinephrine and diphenhydramine (Benadryl)
- Pediatric & adult airways (small, medium, and large)
- Sphygmomanometer (child, adult & extra-large cuffs) and stethoscope
- Pediatric & adult size pocket masks with one-way valve
- Alcohol swabs
- Tongue depressors
- Flashlight with extra batteries (for examination of mouth and throat)
- Wrist watch
- Tourniquet
- Cell phone or access to an on-site phone

(Page 1 of 2)

Emergency Medical Protocol for Management of Anaphylactic Reactions in Children and Teens

Signs and Symptoms of Anaphylactic Reaction

Sudden or gradual onset of generalized itching, erythema (redness), or urticaria (hives); angioedema (swelling of the lips, face, or throat); bronchospasm (wheezing); shortness of breath; shock; abdominal cramping; or cardiovascular collapse.

Treatment in Children and Teens

- If itching and swelling are confined to the injection site where the vaccination was given, observe patient closely for the development of generalized symptoms.
- If symptoms are generalized, activate the emergency medical system (EMS; e.g., call 911) and notify the on-call physician. This should be done by a second person, while the primary nurse assesses the airway, breathing, circulation, and level of consciousness of the patient.
- Administer aqueous epinephrine 1:1000 dilution (i.e., 1 mg/mL) intramuscularly; the standard dose is 0.01 mg/kg body weight, up to 0.3 mg maximum single dose in children and 0.5 mg maximum in adolescents (see chart below).
- In addition, for anaphylaxis, administer diphenhydramine either orally or by intramuscular injection; the standard dose is 1 mg/kg body weight, up to 30 mg maximum dose in children and 100 mg maximum dose in adolescents (see chart below).
- Monitor the patient closely until EMS arrives. Perform cardiopulmonary resuscitation (CPR), if necessary, and maintain airway. Keep patient in supine position (flat on back) unless he or she is having breathing difficulty. If breathing is difficult, patient's head may be elevated, provided blood pressure is adequate to prevent loss of consciousness. If blood pressure is low, elevate legs. Monitor blood pressure and pulse every 5 minutes.
- If EMS has not arrived and symptoms are still present, repeat dose of epinephrine every 10–20 minutes for up to 3 doses, depending on patient's response.
- Record all vital signs, medications administered to the patient, including the time, dosage, response, and the name of the medical personnel who administered the medication, and other relevant clinical information.
- Notify the patient's primary care physician.

Suggested Dosing of Epinephrine and Diphenhydramine

Age Group Dose	Weight * in kg	Weight (lbs)* in lbs	Epinephrine Dose 1 mg/mL injectable (1:1000 dilution) intramuscular	Diphenhydramine (Benadryl) 12.5 mg/5 mL liquid 25 and 50 mg capsules or tabs 50 mg/mL injectable
1–6 mos	4–7 kg	9–15 lbs	0.05 mg (0.05 ml)	5 mg
7–18 mos	7–11 kg	15–24 lbs	0.1 mg (0.1 ml)	10 mg
19–36 mos	11–14 kg	24–31 lbs	0.15 mg (0.15 ml)	15 mg
37–48 mos	14–17 kg	31–37 lbs	0.15 mg (0.15 ml)	20 mg
49–59 mos	17–19 kg	37–42 lbs	0.2 mg (0.2 ml)	
5–7 yrs	19–23 kg	42–51 lbs	0.2 mg (0.2 ml)	30 mg
8–10 yrs	23–35 kg	51–77 lbs	0.3 mg (0.3 ml)	
11–12 yrs	35–45 kg	77–99 lbs	0.4 mg (0.4 ml)	40 mg
13 yrs & older	45+ kg	99+ lbs	0.5 mg (0.5 ml)	50–100 mg

*Dosing by body weight is preferred.

These standing orders for the medical management of vaccine reactions in child and teenage patients shall remain in effect for patients of the _____ until rescinded or until _____.

name of clinic *date*

Medical Director's signature _____ Effective date _____

Sources: American Academy of Pediatrics. Passive Immunization. In: Pickering LK, ed. *Red Book: 2006 Report of the Committee on Infectious Diseases*. 27th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2006: 64–66.
American Pharmacists Association, Grabenstein, JD, *Pharmacy-Based Immunization Delivery*, 2002.

(Page 2 of 2)

www.immunize.org/catg.d/p3082a.pdf • Item #P3082a (8/06)